(W/ Rywichiro Kitano, Rywtaro Matsudo, Norikazu Yamada (KEK) 2010. 08810 & 2102.08784 [hep-lat]

lodoy 4D Pure Yang-Mills Theory W/ A-angle W/ G=SU(2) Q: Free Energy $F(\theta) = -\frac{1}{V} \ln \frac{Z(\theta)}{Z(0)}$ as a function of ΘZ Q: Fate of CP - sym. @ $f = \pi ?$ Q: gapped (confinement)? gapless?

$$\frac{\text{Instanton}(\text{DIGA})[\text{Hooft}]}{\text{F(}\theta\text{I})} = \frac{1}{3}N$$

$$F(\theta) \sim \int_{\rho^{5}}^{\rho \to \infty: \text{IR}} e^{-\frac{8\pi^{2}}{97\omega}}(\mu\rho) \int_{\rho^{5}}^{\mu^{2}} (1 - \cos\theta)$$

$$+ (2 - \text{instanton}) = 1$$



-Xi Works well at TZTC

 \dot{X} . However, not correct for $T \ll Te$ ($p \rightarrow \infty$ divergent : IR problem)

$$\frac{\text{Lorge } N}{\sqrt{2}} \quad [Witten]$$

$$\mathcal{L} \sim \frac{1}{N^{1}} \left(\frac{1}{\sqrt{2}N} \text{Tr } F_{n} + F + \frac{1}{N} \text{Tr } F_{n} F \right)$$

$$\int \int \frac{1}{\sqrt{2}N} \text{Tr } F_{n} + F + \frac{1}{N} \text{Tr } F_{n} F \right)$$

$$\int \int \frac{1}{\sqrt{2}N} \text{Tr } F_{n} + F + \frac{1}{N} \text{Tr } F_{n} F \right)$$

$$\int \int \frac{1}{\sqrt{2}N} \frac{1}{\sqrt{2}N} \frac{1}{\sqrt{2}N} + \frac{1}{\sqrt{2}N} \int \frac{1}{\sqrt{2}N} \frac{1}{\sqrt{2}N} \int \frac{1}{\sqrt{2}N} \frac{1}{\sqrt{2}N} \int \frac{1}{\sqrt{2}N} \frac{1}{\sqrt{2}N} \int \frac{1}{\sqrt{2}$$









generate gauge conf. at
$$\theta = 0$$
 to sign problem
we asure top. charge Q
 $\chi = \frac{\langle Q^2 \rangle_{\theta=0}}{V},$
 $b_2 = -\frac{\langle Q^4 \rangle_{\theta=0} - 3 \langle Q^2 \rangle_{\theta=0}^2}{12 \langle Q^2 \rangle_{\theta=0}},$
 $b_4 = \frac{\langle Q^6 \rangle_{\theta=0} - 15 \langle Q^2 \rangle_{\theta=0} \langle Q^4 \rangle_{\theta=0} + 30 \langle Q^2 \rangle_{\theta=0}^3}{360 \langle Q^2 \rangle_{\theta=0}},$
F(θ) = $5 \chi \theta^2 (1 + b_2 \theta^2 + b_4 \theta^4 + \cdots)$
cf. pure notural in flation (No mura - Wateri - MY (17))
No mura - MY (17))









$$\frac{\chi}{T_c^4} = 0.200(39) , \qquad \frac{\chi^{1/4}}{T_c} = 0.674(31) , \qquad b_2 = -0.049(20) ,$$

seems to be the first determination of $b_2!$
(cf. Bonanno, Bonati, D'Elia (i8) $b_4 = \delta(2) \cdot 10^{-4}$)







Norit~1,5



[Kitano, Matsudo, Yamada, MY (21)] Subvolume Method (cf. [Keith-Hynes, Thacker ('08)] for 2d CP-model) $e^{-V_{sub}}F_{sub}(\theta) = \frac{1}{Z_{loj}}\int \partial U e^{-s_{d}^{2}+i\theta Q_{sub}}$ $= \left\langle e^{i\theta Q sub} \right\rangle = \left(\cos(\theta Q sub) \right)$ $\theta = 0$ F' $F_{sub}(\theta) \sim F(\theta) + \frac{S(\theta)}{l} + O(\frac{1}{l^2})$ Surface tension Inside region QSub & X Zu $(aTc)^{-4} \ll Vsvb \ll Vfull$ better w/ sign problem [

-
$$Su(2)$$
 YM (Symanzik)
- $\beta = \frac{4}{g^2} = 1.975 \xrightarrow{1}{(aT_c)} = 9.50$
- $V_{full} = 24^3 \times \left\{\frac{48}{1}, \frac{8}{1}, 6\right\}_{T=0} = 1.5T_c$

$$- \# (config) = \{68000, 10000, 10000\} (atc) < l < lful
- Vsub = l^4 w) l \in \{10, 12, ..., 24\}$$

$$- lineor extrapolation Fsub(0) = F(0) + \frac{S(0)}{l} + O(\frac{1}{l^2})$$

$$\frac{10000}{resulf}$$

 $f(\theta) \odot T = 1.2 T_c 7 T_c$

f() T=0; clearly NOT 2II-periodic

θ

Summory • $F(\theta)$ for 4d SU(2) YM for $0 \le 0 \le \frac{3\pi}{2}$ using subvolume method (despite sign problem!!)

Summory X4d SU(2) YM: still "large N" spontaneous CP preaking, mass gap $\frac{\chi'^4}{T_c} = 0.674(31) \quad b_2 = -0.049(20)$ (Quantitatively different from 2d CPN-1 model] CP broken CP unbroken Japped Japless NCP & Narit N=1 N=4 N=3 N=2

Future Works

- · Improve systematics
- Explore